
Working Towards a Clinically Usable Allied Health Outcome Set for Paediatric and Adolescent Rheumatology Musculoskeletal Pain Conditions: A Preliminary Scoping Exercise

Victoria Harbottle MSc; BSc (Hons) Physiotherapy [a*]

Ellie Potts BSc (Hons) Physiotherapy [b], Verna Cuthbert BSc (Hons) Physiotherapy; BSc (Hons) [c]

[a] Great North Children's Hospital, Royal Victoria Infirmary, Queen Victoria Road, Newcastle upon Tyne, NE1 4LP

[b] Evelina London Children's Hospital, Guys' and St Thomas' Hospital NHS Trust, Westminster Bridge Road, London, SE1 7EH

[c] Royal Manchester Children's Hospital, Oxford Road, Manchester, M13 9WL

*Corresponding author: victoria.harbottle@ncl.ac.uk

ABSTRACT

Background:

Musculoskeletal conditions are very prevalent within the general population. This is also true for children and young people (CYP), with many seeking care for musculoskeletal pain. Allied Health Professionals (AHPs) are well placed to manage these patients but there is a wide variability in interventions, with no current standardisation in treatment, services, or outcomes.

Aim:

To develop a consensus-based outcome set for clinical use by AHP's working with CYP with musculoskeletal pain conditions

Method:

A staged modified nominal group technique was used. An expert panel of AHP's working in tertiary paediatric rheumatology centres in the UK identified nine domains: fatigue, muscle strength, hand function, school attendance, stamina, balance, sleep, quality of life and goal setting. Literature search and panel discussion preceded anonymous voting to select measures for each domain. Face validity of the final set was tested with a group of YP.

Results:

Consensus was achieved for six of the nine domains. Consensus could not be reached for hand function, sleep, and goal setting in part due a lack of paediatric specific measures, time taken to administer, or cost. YP agreed with the set domains but felt strongly that pain should be included.

Conclusion:

A six domain clinical outcome set has been developed for AHP's treating YP with musculoskeletal pain conditions where time, space, money, and ease of use is paramount. Future work is needed to further develop this set.

Key Points

- 1) Consensus-driven outcome sets for clinical use can be produced with limited funding and may be useful in guiding clinician's outcome selection in practice.
- 2) Young people's views into what is important in allied health professional care may vary from clinicians, and so must be included early in decisions around outcome domains and tools.
- 3) Further work to is needed to consider tools to capture pain symptoms as well as tools in domains that did not reach consensus and to address practical questions around outcome use and frequency in practice.

Introduction

Musculoskeletal (MSK) pain conditions are very prevalent within the general population; evidence suggests that this is also true for children and young people (CYP), with prevalence estimates ranging from 8.5-40% (Tan et al., 2018). It was once thought that MSK pain in CYP was self-limiting and without long-term impact, however, recent work has highlighted the significant impact persisting MSK pain can have on a CYP's quality of life, education, social life, sporting and leisure function and mental health (Scottish Government, 2018, Kamper and Williams, 2017, Liozzi et al., 2019, Pourbordbari et al., 2019). Moreover, there can be longer term

consequences for CYP persisting with pain into adulthood, leading to increased ongoing healthcare use, long-term detriment to health and increased societal burden (Kamper et al., 2016). Significant numbers of CYP are seeking care for MSK pain, within primary, secondary, and tertiary services, and Allied Health Professionals (AHP's), with suitable paediatric experience and training, are well placed to provide interventions for these individuals. At present, however, there is significant variability in both content of and access to evidence-based interventions (Jay and Howard, 2016). Furthermore, there is little guidance on which to base provision, standardise treatment, services, or outcomes.

To address issues of health improvement and reducing health inequalities, we need robust, high quality patient outcome data to work with. Patient outcome data provides crucial clinical information to drive improvements in both the quality of, and decision-making in healthcare. The Chartered Society of Physiotherapy website states that improving the quality of healthcare in the UK is of utmost importance and that clinical outcome measures are one way of providing robust data to drive this improvement (Chartered Society of Physiotherapy, 2020). Despite this, AHP engagement in the use of outcome measurement is variable (Braun et al., 2018). Barriers to the use of outcome measures cited by professionals include time, lack of knowledge in measures to select, and lack of training in their use (Duncan and Murray, 2012).

Consensus derived outcome sets present one way to address challenges with outcome measurement and provide clinicians with an agreed set of standardised outcomes to use in a specified patient group (Williamson et al., 2012). Such sets have historically been developed for clinical trial use, aiming to align outcomes such that comparisons or meta-analyses can be more easily adopted. However, there have been sets developed that aim to guide clinical practice, as well as those aiming to cross both areas, with recognition that sets for clinical use provide their own challenges (Dodd et al., 2020).

Within paediatric and adolescent rheumatology recent advances have included developments for a minimum clinical data set for Juvenile Idiopathic Arthritis; CAPTURE (McErlane et al., 2020) and Juvenile Dermatomyositis (McCann et al., 2015). These sets are consensus driven and specify outcomes or measures which reflect current clinical practice and include both clinician-reported and patient/parent reported outcome measures.

The work reported here was initially undertaken by the British Society for Rheumatology; Paediatric and Adolescent Rheumatology (BSpaR) AHP group, who recognised the need to consider outcome measures which inform clinical practice undertaken by AHPs specifically, and to generate a set reflecting the challenges and needs of AHPs working in the UK. To our knowledge this was the first consensus driven outcome set developed for AHP use in this area. As such we aimed to scope a clinical set of outcome tools to guide AHP's working with CYP with MSK pain conditions, and as a starting point for a future research driven core outcome set. Our core objectives were:

- (i) Generate a set of consensus-derived outcome measures that are widely accessible and suitable for use in a clinical setting to guide clinician use in MSK settings.
- (ii) Sense-check this set within a group of service users.
- (iii) Make recommendations for further work developing a robust minimal core outcome set suitable for routine patient data collection and clinical research.

Method

A multi-phased methodological approach was used, as shown in Figure 1:




Figure 1: An overview of the multi-staged methodology used.

Phase 1: Current AHP members of BSpaR, working as senior clinicians (at least 5 years' experience specialising in paediatric rheumatology) were contacted by email and invited to participate as panel members. Fourteen clinicians, detailed in table 1, agreed to take part in this work, which was not funded and was carried out alongside usual clinical workload.

Table 1: Professions and centres of expert panel members

Profession	Physiotherapists	7
	Occupational Therapists	6
	Psychologists	1
Centre	Alder Hey Children's Hospital	1
	Birmingham Children's Hospital	2
	Bristol Royal Hospital for Children	1
	Evelina London Children's Hospital	4
	Great Ormond Street Hospital	2
	Great North Children's Hospital	1
	Royal Manchester Children's Hospital	2
	Southampton Children's Hospital	1

Initial roundtable discussion focused on the aims of the piece of work and agreeing on what domains were important for AHP's to measure. Anonymous voting on the domains listed by post-it notes was carried out by the expert AHP panel and results were used to identify the priority domains addressed in Phase 2.

Phase 2: The domains identified were shared out within the panel. An informal review of the literature was carried out for each of the outcome domains selected, to identify potential outcome measures which had been or were currently being used within paediatric musculoskeletal or rheumatology clinical settings and research studies. Electronic databases PubMed, CINAHL, Medline and web of science were systematically searched. Search terms included the domain name AND rheumatology OR musculoskeletal. Searches were limited to studies involving children, and young people, published in English, and from 2010 onwards, to limit focus to the most recent measures. Abstracts were used where possible to identify outcome measures, with full text sought if this was not possible. Additionally, any measures currently used by panel members or colleagues in clinical practice with the defined population were included.

Phase 3: A meeting of the panel was convened. Each domain and associated literature findings were presented to the panel with discussion facilitated by chairperson to focus on the advantages and disadvantages of each

measure. Discussion was guided by the principles for outcome tool selection suggested by Prinsen et al (Prinsen et al., 2016) with particular focus on:

- Validity within a paediatric and adolescent population
- Use in a variety of clinical settings
- Cost to obtain
- Ease of use with minimal training

Phase 4: Following the face-to-face panel meeting, a survey, using survey monkey (www.surveymonkey.com), was sent to each of the panel members by email, asking the participants to choose the strongest measure for each domain, measured against the list above, or none, if none of the measures were felt to fulfil the criteria. Consensus was pre-determined as agreement between experts, participating in the online survey, of 60% or more (Fink et al., 1984).

Phase 5: The consensus findings were sense checked by members of the national youth advisory panel in adolescent and young adult rheumatology – Your Rheum (<http://yourrheum.org>; (McDonagh et al., 2019)). Meetings with young people were conducted without presence of clinicians to reduce potential influence by health professionals and to optimise the willingness of the young people to input authentically to the discussion. The meeting involved three young people aged 12, 18 and 24. An arts-based approach was selected for this event in order to generate thoughts, ideas and discussion about the topic prior to the use of open-ended questions about the COS. Art-based approaches are often used in eliciting YP views around health-related issues and are frequently used in conjunction with interviews or discussion groups (Coad, 2007). Table 2 shows the methods used for the face validity testing.

Table 2: Arts-based face validity testing method

Step One	Young people (YP) were given a size A0 “life course map” and a variety of craft materials: pens, pencils, magazines, scissors, glue etc. Instruction was given to consider broadly the outcomes, their indicators of health and well-being and key events throughout childhood/adolescence and young adulthood that were important and had meaning to them.
Step Two	YP were asked to record outcomes, in any medium available, at the point on the life course map where they felt they fitted best. It was impressed upon the YP that there was no right or wrong in this situation and that they were free to record and express anything that they felt was of importance within this topic area.
Step Three	Following this the outcomes identified by the expert AHP group were introduced to the YP. They were invited to comment openly on the domains, specifically: • Did they agree or disagree with the items included? • Was there anything that had not been included that should have? YP feedback was documented during the discussion and used to inform the final COS.

Results

Phase 1 findings: Nine domains were agreed on. These were fatigue, sleep, school function, achievement of goals, muscle strength, balance, stamina, hand function and quality of life.

Phase 2 findings: Literature scoping identified 39 potential measures across the 9 domains (See appendix 1).

Phase 3 and 4 findings: The response rate for the expert panel survey was 83% (n=10 of 12). Consensus was derived for 6 of the 9 domains considered, with no consensus for sleep, achievement of goals or hand function. Table 3 shows the agreed core outcome set. Fatigue, muscle strength and stamina produced a tied result and so both measures have been included at this stage.

Table 3: List of agreed measures for core outcome set

Domain	Outcome Measure
Fatigue	Fatigue Visual Analogue Scale (Crawford et al., 2011), Paediatric Quality of Life Multidimensional Fatigue Scale (Varni et al., 2004)
School function	Attendance rate (Weitzman, 1986)
Muscle Strength	Kendal scale (Kendall et al., 1993), Oxford scale (Compston, 1942)
Balance	Standardised Single Leg Stance (Condon and Cremin, 2014)
Stamina	Timed step test (Balfour-Lynn et al., 1998), Two minute walk test (2MWT) (Bohannon et al., 2014)
Quality of life	Paediatric Quality of Life Inventory Generic Core Measure (Peds QL) (Varni et al., 1999)

Phase 5 findings: Discussion with YP found agreement with all domains included in the COS, and they felt that they were all of importance to their daily lives and functioning. However, they felt very strongly that pain was a domain that should be included due to the large impact it has on their day-to-day life.

Discussion

A preliminary core outcome set for clinical use by allied health professionals working with children and young people with musculoskeletal pain conditions has been developed, comprising of six consensus agreed outcome measures for fatigue, school function, muscle strength, stamina, balance, and quality of life. Consensus could not be reached for measures for sleep, goal setting and hand function. Possible reasons for this include identified tools not having suitable psychometric properties, being expensive to obtain, being time consuming to use in clinical practice and lacking appropriate validity testing.

Core outcome sets are an accepted way of agreeing a standardised minimum set of outcome measures (Williamson et al, 2012). Interest in the development of such sets has resulted in a series of published methodological guidance The COS-STAD checklist for outcome selection, along with guidance on reporting, protocol publishing and outcome tool selection (Kirkham et al., 2017, Kirkham et al., 2016, Kirkham et al., 2019, Prinsen et al., 2016) is available to guide researchers. The methodology used here has several strengths, including a clear setting, population, condition, and interventions, allowing clinicians to judge the suitability of the set for their use. However, there are several limitations. Unlike traditional COS development, both domain selection and outcome tool selection were encompassed within the one multi-phased study. Full COS development places more emphasis on psychometric evaluation rather than feasibility of use, setting a minimum standard for inclusion of an outcome measure (Prinsen et al., 2016). Barriers to outcome measure use for AHPs highlight lack of time as a key factor in uptake (Duncan and Murray, 2012). Therefore, we made the decision at this early stage to prioritise feasibility over psychometric properties, agreeing validation in a paediatric population, as minimum criterion. A further limitation was in stakeholder inclusion. The stakeholder group included representation from eight centres and three professional groups but lacked input from either CYP or parents. Patient involvement is noted as important in COS development (Kirkham et al., 2017, Williamson et al., 2012). We were able to seek the opinions of CYP through the research involvement group Your Rheum (www.yourrheum.org). Due to resources this was limited to a retrospective sense checking activity. This was attended by three YP, aged between twelve and twenty-four, with no inclusion of parents or younger children. As this activity was conducted through the Patient and Public Involvement (PPI) framework and organised by Your Rheum, this limited the research groups' input into the number of participants involved. We recognise the limitations of this small sample size, and the associated impact of limiting the generalisability of findings to a wider population. Additionally, the narrow age range restricts the insight gained to the adolescent population, with additional work needed to consider use of the set in a younger population.

As the first allied health clinical core outcome set in paediatric musculoskeletal conditions it is not possible to directly compare our findings to other studies. Other, comparable work in progress includes an arthritis symptom tracker for teens with JIA. This project has sought the opinions of young people with JIA on which items and symptoms they felt were most important to monitor. YP identified energy levels, medication side effects, activity levels and sleep function as important to track the course of their disease (Versus Arthritis, 2020). These map well to the outcomes of fatigue, stamina and strength, and sleep identified in this study, with medication side effects not directly within allied health professionals' scope of practice. However, there were areas of disagreement between this project and our core outcome set.

When considering domain inclusion, pain was excluded by the professional panel in the early stages. This was because the panel prioritised functional outcomes, as it was felt that focus on controlling pain may lead to prioritisation of pain symptom management over functional improvement. This has been found to be unhelpful in longer term instances of pain, and can result in dysfunctional beliefs by young people and their parents around pain (Caes et al., 2012). Additionally, it was felt that the known complexity of pain assessment, as well as the complex interaction of pain and developmental stage would pose a challenge outside the scope of this work. The YP involved in our sense-checking strongly disagreed with the panel, which aligns with findings from the Versus Arthritis app project, where pain was a symptom prioritised by YP to be tracked (Versus Arthritis, 2020). The YP we consulted were clear that pain was central to their function, was needed for interpretation of other results and that it should be personalised to the young person and measured specifying context.

Parents and YP often differ in their treatment priorities to professionals (Sinha et al., 2012, Sherratt et al., 2020), as has been found to be the case in this work. Whilst the panels concern around pain behaviour may be relevant, ensuring that clinical outcomes used are important and meaningful to YP is crucial in ensuring a shared focus for treatment. Failure to capture outcomes deemed relevant to YP may limit the use of clinical decision making, and impair engagement and trust between YP and their clinicians (Sherratt et al., 2020). Whilst the PedsQL generic core measure, recommended as the quality-of-life tool, does include items that related to pain symptoms, pain should be looked at as a further outcome domain. Future work is needed to agree on an appropriate pain outcome tool for use in practice and this work should prioritise engagement of young people early in the research process.

This preliminary piece of work is a valuable starting point for aiding clinical selection of outcome measures. Along with the addition of pain, future work should explore further the measures with tied consensus to allow the recommendation of one measure per domain (Prinsen et al., 2016). Additionally, consideration should be given to those domain tools that did not reach the pre-agreed consensus- sleep, hand function and achievement of goals. Psychological outcomes should also be considered, as the interplay between physical and psychological health and wellbeing has long been recognised. Psychologists have an important role to play in the management of paediatric MSK conditions (Davis et al., 2017), although access is variable nationally (British Society for Rheumatology, 2019, Hawley et al., 2018).

Advice on timing and frequency of outcome collection is also important to consider. AHP models of care vary from blocks of more intense treatment for some patients, to regular review throughout childhood and adolescence to account for developmental changes and to provide developmentally appropriate care. ARMA (Arthritis and Musculoskeletal Alliance) standards for CYP with JIA for example, suggest a minimum of an MDT review annually (Arthritis and Musculoskeletal Alliance, 2010), however this is often at odds with current NHS policies, with patients with long term conditions being discharged from treatment. AHP care models for paediatric MSK conditions vary nationally and will impact agreement on the timing and frequency of COS collection.

Conclusion

This scoping work represents, to the best of our knowledge, the first allied health specific outcome set led by allied health professionals for their clinical use in Paediatric Rheumatology settings. We recommend an initial consensus derived set of outcome measures to assist allied health professionals in the selection of tools for assessing outcomes in young people with musculoskeletal pain conditions. We recognise the limitations to this work and identify key recommendations points for future projects to further develop this core outcome set.

This is an important collaborative step by the BSpaR AHP group to challenge variation in paediatric MSK service provision and work towards driving improvements in care and informing decision-making for this population.

Ethical Approval:

Ethical approval was not sought for this study: attendance by AHP's at the expert panel meeting was considered as consenting to participate in the consensus process. Consent by CYP at the face validity testing was sought by YOURRHEUM personnel and was done within the PPI framework. As such the young people were not research participants and therefore ethics approval was not required (National Research Ethics Service and INVOLVE, 2009).

Declaration of Interests: The authors declare no conflict of interest.

Funding Sources: This research did not receive any specific grant from funding agencies in the public, commercial, or not-for-profit sectors.

Acknowledgements: The authors wish to thank expert panel members, with special thanks to Louise Fox, for their contribution to this work, Dr Janet McDonagh and Dr Lis Cordingley for initial review and feedback, as well as Dr Joyce Davidson, Dr Kate Armon and Dr Flora McErlane for their support and assistance.

References

- Arthritis and Musculoskeletal Alliance (2010). 'ARMA Standards of Care for Children and Young People with Juvenile Idiopathic Arthritis'. Available: www.arma.uk.net. [Accessed 20th December 2019].
- Balfour-Lynn, I M, Prasad, S A, Laverty, A, Whitehead, B F & Dinwiddie, R (1998). 'A step in the right direction: Assessing exercise tolerance in cystic fibrosis'. *Pediatric Pulmonology*, 25, 278-284.
- Bohannon, R W, Bubela, D, Magasi, S, McCreath, H, Wang, Y-C, Reuben, D, Rymer, W Z & Gershon, R (2014). 'Comparison of walking performance over the first 2 minutes and the full 6 minutes of the Six-Minute Walk Test'. *BMC Research Notes*, 7, 269-269.
- Braun, T, Rieckmann, A, Weber, F & Gruneberg, C (2018). 'Current use of measurement instruments by physiotherapists working in Germany: a cross-sectional online survey'. *BMC Health Serv Res*, 18, 810.
- British Society for Rheumatology (2019). 'Paediatric and Adolescent Rheumatology: The State of Play Report '. Available: <https://www.rheumatology.org.uk/News-Policy/Details/Paediatric-Adolescent-Rheumatology-State-of-Play-report> [Accessed 23rd December 2019].
- Caes, L, Vervoort, T, Eccleston, C & Goubert, L (2012). 'Parents who catastrophize about their child's pain prioritize attempts to control pain'. *Pain*, 153, 1695-1701.
- Chartered Society of Physiotherapy (2020). 'Outcome and experience measures '. Available: <https://www.csp.org.uk/professional-clinical/research-evaluation/outcome-experience-measures> [Accessed 15th January 2020].
- Coad, J (2007). 'Using art-based techniques in engaging children and young people in health care consultations and/or research'. *Journal of Research in Nursing*, 12, 487-497.
- Compston, A (1942). 'Aids to the investigation of peripheral nerve injuries'. Medical Research Council: Nerve Injuries Research Committee. His Majesty's Stationery Office, 48-48.
- Condon, C & Cremin, K (2014). 'Static Balance Norms in Children'. *Physiotherapy Research International*, 19, 1-7.
- Crawford, B K, Piault, E C, Lai, C & Bennett, R M (2011). 'Assessing fibromyalgia-related fatigue: content validity and psychometric performance of the Fatigue Visual Analog Scale in adult patients with fibromyalgia'. *Clin Exp Rheumatol*, 29, S34-43.
- Davis, A M, Rubinstein, T B, Rodriguez, M & Knight, A M (2017). 'Mental health care for youth with rheumatologic diseases - bridging the gap'. *Paediatric Rheumatology*, 15.
- Duncan, E A S & Murray, J (2012). 'The barriers and facilitators to routine outcome measurement by allied health professionals in practice: A systematic review'. *BMC Health Services Research*, 12.
- Fink, A, Kosecoff, J, Chassin, M & Brook, R H (1984). 'Consensus methods: Characteristics and guidelines for use'. *American Journal of Public Health*, 74, 979-983.
- Hawley, D P, Howsley, P, McDonagh, J E, McErlane, F, Rapley, T, Tattersall, R S & Webb, N (2018). 'P24 United Kingdom survey of access to psychology services for adolescents and young adults with rheumatic and musculoskeletal disease'. *Rheumatology*, 57, key273-026.
- Jay, M A & Howard, R F (2016). 'Inequalities in access to a tertiary children's chronic pain service: a cross-sectional study'. *Archives of Disability in Childhood*, 101, 657-661.
- Kamper, S J, Henschke, N, Hestbaek, L, Dunn, K M & Williams, C M (2016). 'Musculoskeletal pain in children and adolescents'. *Brazilian Journal of Physical Therapy*, 20, 275-284.
- Kamper, S J & Williams, C M 2017. Musculoskeletal pain in children and adolescents: A way forward. Movement Science Media.

Kendall, F P, McCreary, E K, Provance, P G, Rodgers, M & Romani, W A (1993). *Muscles, testing and function: with posture and pain*, Williams & Wilkins Baltimore, MD.

Kirkham, J J, Davis, K, Altman, D G, Blazeby, J M, Clarke, M, Tunis, S & Williamson, P R (2017). 'Core Outcome Set-STANDards for Development: The COS-STAD recommendations'. *PLoS Medicine*, 14, 1-10.

Kirkham, J J, Gorst, S, Altman, D G, Blazeby, J M, Clarke, M, Devane, D, Gargon, E, Moher, D, Schmitt, J, Tugwell, P, Tunis, S & Williamson, P R (2016). 'Core Outcome Set-STANDards for Reporting: The COS-STAR Statement'. *PLoS Medicine*, 13, 1-11.

Kirkham, J J, Gorst, S, Altman, D G, Blazeby, J M, Clarke, M, Tunis, S & Williamson, P R (2019). 'Core Outcome Set-STANDARDised Protocol Items: The COS-STAP Statement'. *Trials*, 20, 1-7.

Lioffi, C, Johnstone, L, Lilley, S, Caes, L, Williams, G & Schoth, D E (2019). 'Effectiveness of interdisciplinary interventions in paediatric chronic pain management: a systematic review and subset meta-analysis'. *British Journal of Anaesthesia*, 123, e359-e371.

McCann, L J, Kirkham, J J, Wedderburn, L R, Pilkington, C, Huber, A M, Ravelli, A, Appelbe, D, Williamson, P R & Beresford, M W (2015). 'Development of an internationally agreed minimal dataset for juvenile dermatomyositis (JDM) for clinical and research use'. *Trials*, 16.

McDonagh, J E, Lunt, L, Cresswell, K, Parsons, S & Tattersall, R (2019). 'Research priority setting by young people with rheumatic musculoskeletal disease'. *Archives of Disease in Childhood*, 104, 204-204.

McErlane, F, Armit, G, Cobb, J, Bailey, K, Cleary, G, Douglas, S, Lunt, L, Rashid, A, Sampath, S, Shoop-Worrall, S, Smith, N, Foster, H & Thomson, W (2020). 'CAPTURE-JIA: a consensus-derived core dataset to improve clinical care for children and young people with juvenile idiopathic arthritis'. *Rheumatology*, 59, 137-145.

National Research Ethics Service and INVOLVE (2009). 'Patient and public involvement in research and research ethics committee review'. Available: <https://www.invo.org.uk/wp-content/uploads/2011/12/INVOLVENRESfinalStatement310309.pdf> [Accessed 1st March 2021].

Pourbordbari, N, Riis, A, Jensen, M B, Olesen, J L & Rathleff, M S (2019). 'Poor prognosis of child and adolescent musculoskeletal pain: A systematic literature review'. *BMJ Open*, 9, e024921.

Prinsen, C A C, Vohra, S, Rose, M R, Boers, M, Tugwell, P, Clarke, M, Williamson, P R & Terwee, C B (2016). 'How to select outcome measurement instruments for outcomes included in a "Core Outcome Set" - a practical guideline'. *Trials*, 17, 1-10.

Scottish Government (2018). 'Management of chronic pain in children and young people: summary '. Available: <https://www.gov.scot/publications/management-chronic-pain-children-young-people/> [Accessed 15th January 2020].

Sherratt, F C, Bagley, H, Stones, S R, Preston, J, Hall, N J, Gorst, S L & Young, B (2020). 'Ensuring young voices are heard in core outcome set development: international workshops with 70 children and young people'. *Res Involv Engagem*, 6, 19.

Sinha, I P, Gallagher, R, Williamson, P R & Smyth, R L (2012). 'Development of a core outcome set for clinical trials in childhood asthma: a survey of clinicians, parents, and young people'. *Trials*, 13, 103.

Tan, A, Strauss, V Y, Protheroe, J & Dunn, K M (2018). 'Epidemiology of paediatric presentations with musculoskeletal problems in primary care'. *BMC Musculoskeletal Disorders*, 19, 1-6.

Varni, J W, Burwinkle, T M & Szer, I S (2004). 'The PedsQL™ multidimensional fatigue scale in pediatric rheumatology: Reliability and validity'. *Journal of Rheumatology*, 31, 2494-2500.

Varni, J W, Seid, M & Rode, C A (1999). 'The PedsQL™: Measurement Model for the Pediatric Quality of Life Inventory'. *Medical Care*, 37, 126-139.

Versus Arthritis. (2020). Available: <https://www.versusarthritis.org/about-arthritis/young-people/arthritis-tracker/> [Accessed 10th January 2020].

Weitzman, M (1986). 'School absence rates as outcome measures in studies of children with chronic illness'. *Journal of Chronic Diseases*, 39, 799-808.

Williamson, P R, Altman, D G, Blazeby, J M, Clarke, M, Devane, D, Gargon, E & Tugwell, P (2012). 'Developing core outcome sets for clinical trials: Issues to consider'. *Trials*, 13, 1-8.

Appendix 1: All outcome measures identified and considered by expert panel

Domain	Measures considered	Reference
Fatigue	Fatigue Visual Analogue Scale (Fatigue VAS)	Crawford BK, Piau EC, Lai C, Bennett RM. Assessing fibromyalgia-related fatigue: content validity and psychometric performance of the Fatigue Visual Analog Scale in adult patients with fibromyalgia. <i>Clinical and experimental rheumatology</i> . 2011;29(6 Suppl 69):S34-43.
	Paediatric Quality of Life Inventory (Peds QL) Multidimensional Fatigue Scale	Varni JW, Burwinkle TM, Szer IS. The PedsQL™ multidimensional fatigue scale in pediatric rheumatology: Reliability and validity. <i>Journal of Rheumatology</i> . 2004;31(12):2494-500.
	Computerised Adaptive Test for Fatigue in Rheumatoid Arthritis (CAT Fatigue RA)	Nikolaus S, Bode C, Taal E, Vonkeman HE, Glas CAW, van de Laar MAFJ. Construct Validation of a Multidimensional Computerized Adaptive Test for Fatigue in Rheumatoid Arthritis. <i>PLOS ONE</i> . 2015;10(12):e0145008-e.
	Bristol Rheumatoid Arthritis Fatigue Scales (BRAf)	Nicklin J, Cramp F, Kirwan J, Greenwood R, Urban M, Hewlett S. Measuring fatigue in rheumatoid arthritis: A cross-sectional study to evaluate the Bristol Rheumatoid Arthritis Fatigue Multi-Dimensional questionnaire, visual analog scales, and numerical rating scales. <i>Arthritis Care & Research</i> . 2010;62(11):1559-68.
Sleep	Multidimensional Fatigue Symptom Inventory-Short Form (MFSI-SF)	Stein KD, Jacobsen P, Thors C Further validation of the multidimensional fatigue symptom inventory-short form. <i>Journal of pain and symptom management</i> . 2004;27(1):14-23 .
	Children's Sleep Habits Questionnaire (CSHQ)	Owens JA, Spirito A, McGuinn M. The Children's Sleep Habits Questionnaire (CSHQ): Psychometric Properties of A Survey Instrument for School-Aged Children. <i>Sleep</i> . 2000;23(8):1-9.
	Bedtime problems, Excessive daytime sleepiness, Awakenings during the night, Regularity and duration of sleep, Snoring (BEARS)	Owens JA, Dalzell V. Use of the 'BEARS' sleep screening tool in a pediatric residents' continuity clinic: A pilot study. <i>Sleep Medicine</i> . 2005.
School Function	Attendance Rate	Weitzman M. School absence rates as outcome measures in studies of children with chronic illness. <i>Journal of Chronic Diseases</i> . 1986;39(10):799-808.
	School Setting Interview (SSI)	Hemmingsson H, Borell L. The Development of an Assessment of Adjustment Needs in the School Setting for Use with Physically Disabled Students. <i>Scandinavian Journal of Occupational Therapy</i> . 1996;3(4):156-62.

	School Function Assessment (SFA)	Coster WJ, Mancini MC, Ludlow LH. Factor Structure of the School Function Assessment. Educational and Psychological Measurement. 1999;59(4):665-77.
Achievement of Goals	Goal Attainment Scale (GAS)	Steenbeek D, Ketelaar M, Galama K, Gorter. JW. Goal attainment scaling in paediatric rehabilitation: a critical review of the literature. Developmental Medicine & Child Neurology. 2007;49(7):550-556.
	Canadian Occupational Performance Measure (CPOM)	Cusick A, Lannin N, Lowe K. Adapting the Canadian Occupational Performance Measure for use in Paediatric clinical trials. J Disab and Rehab. 2007, 29(10):761.
	Child and Adolescent Mental Health Service Outcomes Research Consortium/ Improving Access to Psychological Therapies Goal Based Outcome Measure (CORC/IAPT GBO)	Wolpert M, Ford T, Trustam E, Law D, Deighton J, Flannery H, et al. Patient-reported outcomes in child and adolescent mental health services (CAMHS): Use of idiographic and standardized measures. Journal of Mental Health. 2012;21(2):165-73.
Muscle Strength	Grip Strength	Hamam N, Abdel-Wahab N, Rashed AM. FRI0552 Hand Grip Strength as Simple Predictor of Disease Activity and Disability in Patients with Juvenile Idiopathic Arthritis. Annals of the Rheumatic Diseases. 2014;73(Suppl 2):586.
	Kendall Scale	Kendall FP, McCreary EK, Provance PG, Rodgers M, Romani WA. Muscles, testing and function: with posture and pain: Williams & Wilkins Baltimore, MD; 1993.
	Oxford Scale	Compston A. Aids to the investigation of peripheral nerve injuries. Medical Research Council: Nerve Injuries Research Committee His Majesty's Stationery Office. 1942:48.

Balance	Star Excursion Balance Test (SEBT)	Hertel J, Miller SJ, Denegar CR. Intratester and Intertester Reliability during the Star Excursion Balance Tests. <i>Journal of Sport Rehabilitation</i> . 2000;9(2):104-16.
	Movement Assessment Battery for Children (Movement ABC)	Henderson SE, Sugden DA, Barnett AL. Movement assessment battery for children-2 second edition [Movement ABC-2]. London, UK: The Psychological Corporation. 2007.
	The Bruininks–Oseretsky test of motor proficiency- second edition (BOT-2)	Bruininks RH, Bruininks BD. BOT2: Bruininks-Oseretsky test of motor proficiency: AGS Publishing; 2005.
	Timed Up and Go (TUG)	Williams EN, Carroll SG, Reddihough DS, Philips BA, Galea MP. Investigation of the timed 'Up & Go' test in children. <i>Developmental Medicine and Child Neurology</i> . 2005;47(8):518-24.
	Standardised Single Leg Stance (Standardised SLS)	Condon C, Cremin K. Static Balance Norms in Children. <i>Physiotherapy Research International</i> . 2014;19(1):1-7.
Stamina	Six Minute Walk Test (6MWT)	Enright PL. The six-minute walk test. <i>Respiratory care</i> . 2003;48(8):783-5.
	Two Minute Walk Test (2MWT)	Bohannon RW, Bubela D, Magasi S, McCreath H, Wang YC, Reuben D, et al. Comparison of walking performance over the first 2 minutes and the full 6 minutes of the Six-Minute Walk Test. <i>BMC Research Notes</i> . 2014;7(1).
	Five Times Sit-To-Stand	Kumban W, Amatachaya S, Emasithi A, Siritaratiwat W. Five-times-sit-to-stand test in children with cerebral palsy: Reliability and concurrent validity. <i>NeuroRehabilitation</i> . 2013;32(1):9-15.
	Timed Step Test	Balfour-Lynn IM, Prasad SA, Lavery A, Whitehead BF, Dinwiddie R. A step in the right direction: Assessing exercise tolerance in cystic fibrosis. <i>Pediatric Pulmonology</i> . 1998;25(4):278-84.

Hand Function	Cochin Hand Function (COCHIN)	Duruöz MT, Poiraudeau S, Fermanian J, Menkes CJ, Amor B, Dougados M, et al. Development and validation of a rheumatoid hand functional disability scale that assesses functional handicap. <i>Journal of Rheumatology</i> . 1996;23(7):1167-72.
	Juvenile Arthritis Functional Assessment Scale (JAFAS)	Lovell DJ, Shear E, Hartner S, McGirr G, Schulte M, Levinson J, et al. Development of a disability measurement tool for juvenile rheumatoid arthritis. The juvenile arthritis functional assessment scale. <i>Arthritis & Rheumatism</i> . 1989;32(11):1390-5.
	Arthritis Hand Function Test (AHFT)	Backman C, Mackie H, Harris J. Arthritis Hand Function Test: Development of a Standardized Assessment Tool. <i>The Occupational Therapy Journal of Research</i> . 1991;11(4):245-56.
	Grip Ability Test (GAT)	Dellhag B, Bjelle A. A Grip Ability Test for use in rheumatology practice. <i>The Journal of rheumatology</i> . 1995;22(8):1559-65.
	Functional Index for Hand Osteoarthritis (FIHOA)	Dreiser R-L, Maheu E, Guillou GB, Caspard H, Grouin J-M. Validation of an algofunctional index for osteoarthritis of the hand. <i>Revue du Rhumatisme et des Maladies Osteoarticulaires-Edition Francaise</i> . 1995;62(6):43S-S.
	Jebsen Hand Function Test (JHFT)	Jebsen RH, Taylor N, Trieschmann RB, Trotter MJ, Howard LA. An objective and standardized test of hand function. <i>Archives of physical medicine and rehabilitation</i> . 1969;50(6):311-9..
	Michigan Hand Outcomes Questionnaire (MHQ)	Chung KC, Pillsbury MS, Walters MR, Hayward RA. Reliability and validity testing of the Michigan Hand Outcomes Questionnaire. <i>Journal of Hand Surgery</i> . 1998;23(4):575-87.
	Australian/Canadian Osteoarthritis Hand Index (AUSCAN)	Bellamy N, Campbell J, Haraoui B, Buchbinder R, Hobby K, Roth JH, et al. Dimensionality and clinical importance of pain and disability in hand osteoarthritis: Development of the Australian/ Canadian (AUSCAN) osteoarthritis hand index. <i>Osteoarthritis and Cartilage</i> . 2002;10(11):855-62.
	Pizza Putty	Leese S, Hackett J, Aslam R, Mills L, Jester A, Southwood TR. A91: The Development and Initial Evaluation of a Standardised Play Based Hand Assessment Tool for Use in Children and Adolescents: the PizzaPutty™ Test. <i>Arthritis & Rheumatology</i> . 2014;66:S126-S.

Quality of Life	Paediatric Quality of Life Inventory Generic Core Measure (Peds QL (Generic Core))	Varni JW, Seid M, Rode CA. The PedsQL™: Measurement Model for the Pediatric Quality of Life Inventory. 1999.
	Childhood Health Assessment Questionnaire (CHAQ)	Nugent J, Ruperto N, Grainger J, Machado C, Sawhney S, Baildam E, et al. The British version of the childhood health assessment questionnaire (CHAQ) and the child health questionnaire (CHQ). Clinical and experimental rheumatology. 2001;19(4; SUPP/23):S163-S7.
	Child Health Questionnaire (CHQ)	Nugent J, Ruperto N, Grainger J, Machado C, Sawhney S, Baildam E, et al. The British version of the childhood health assessment questionnaire (CHAQ) and the child health questionnaire (CHQ). Clinical and experimental rheumatology. 2001;19(4; SUPP/23):S163-S7.
	Musculoskeletal Health Questionnaire (MSK-HQ)	Hill JC, Kang S, Benedetto E, Myers H, Blackburn S, Smith S, et al. Development and initial cohort validation of the Arthritis Research UK Musculoskeletal Health Questionnaire (MSK-HQ) for use across musculoskeletal care pathways. BMJ Open. 2016;6(8).